Application No. 10/579,357

Attorney Docket No.: 06478.1507-00000

## **AMENDMENTS TO THE CLAIMS:**

This listing of claims will replace all prior versions and listings of claims in the application:

1. (Currently Amended) A stable <u>proteinimmunoglubulin</u> preparation, wherein the preparation comprises one or more stabilisers selected from the group-consisting of non-polar and basic amino acids<u>proline</u> and wherein the preparation has a pH of 4.2 to 5.4, and wherein the preparation does not comprise nicotinamide.

## 2-3. (Cancelled)

- 4. (Currently Amended) The preparation of claim 31, wherein proline is L-proline.
- 5. (Previously presented) The preparation of claim 1, wherein said preparation has a pH of 4.5 to 5.2.
- 6. (Previously presented) The preparation of claim 5, wherein said preparation has a pH of 4.6 to 5.0.
- 7. (Currently amended) The preparation of claim 1, wherein said preparation comprises the one or more stabilisers at athe final concentration of proline is at least 0.2 M.

Application No. 10/579,357

Attorney Docket No.: 06478.1507-00000

8. (Currently amended) The preparation of claim 7, A stable immunoglobulin preparation, wherein said preparation comprises the one or more stabilisers at a proline and has a pH of 4.2 to 5.4, and wherein the final concentration of proline is between 0.2 to 0.4 M.

- 9. (Currently amended) The preparation of claim 1 or 8, wherein saidpreparation comprises the one or more stabilisers at athe final concentration of proline
  is 0.25 M.
- 10. (Currently amended) The preparation of claim 1 or 8, wherein the protein immunoglobulin concentration of said preparation is from 5 to 25% w/v.
- 11. (Currently amended) The preparation of claim 10, wherein the proteinimmunoglobulin concentration of said preparation is from 15 to 20% w/v for subcutaneous administration.
- 12. (Currently amended) The preparation of claim 10, wherein the preteinimmunoglobulin concentration of said preparation is from 6 to 15% w/v, for intravenous administration.
- 13. (Currently amended) The preparation of claim 12, wherein the proteinimmunoglobulin concentration of said preparation is from 8 to 12% w/v.

Application No. 10/579,357 Attorney Docket No.: 06478.1507-00000

14. (Cancelled)

15. (Currently amended) The preparation of claim 44<u>1 or 8</u>, wherein said preparation is an IgG, IgA or IgM preparation.

16. (Currently amended) A pharmaceutical composition comprising the proteinimmunoglobulin preparation of claim 14<u>1 or 8</u> and pharmaceutically acceptable additives.

17. (Cancelled)

18. (Withdrawn, currently amended) A method of stabilising proteinimmunoglobulin preparations, comprising providing an aqueous proteinimmunoglobulin solution and adding one or more stabilisers selected from the group consisting of basic and non-polar amino acidsproline, wherein the pH of the solution is adjusted to a pH of about 4.2 to 5.4, and wherein the preparation does not comprise nicotinamide.

19. (Cancelled)

20. (Withdrawn) The method of claim 18, wherein the pH is adjusted to 4.8.

Application No. 10/579,357

Attorney Docket No.: 06478.1507-00000

21. (Withdrawn) The method of claim 18, wherein the final concentration of the one or more stabilisers proline is adjusted to between 0.2 to 0.4 M.

## 22. (Cancelled)

- 23. (Currently amended) A pharmaceutical composition comprising the proteinimmunoglobulin preparation of claim 1 and pharmaceutically acceptable additives.
- 24. (New) A method of decreasing aggregate formation and/or of decreasing colouring of immunoglobulin preparations, comprising providing an aqueous immunoglobulin solution and adding one or more stabilisers chosen from non-polar amino acids, wherein the pH of the solution is adjusted to a pH of about 4.2 to 5.4.
  - 25. (New) The method of claim 25, wherein the pH is adjusted to 4.8.
- 26. (New) The method of claim 25, wherein the non-polar amino acid is proline.
- 27. (New) The method of claim 26, wherein the proline concentration is adjusted to between 0.2 to 0.4 M.
- 28. (New) The preparation of claim 1 or 8, wherein the final concentration of proline is between 0.2 to 0.3 M.